

Conference Abstract

A short self-paced submaximal test to monitor endurance cycling fitness

Jules Cusson-Fradet ^{1,2,*}, Frédéric Domingue ^{2,3} and Claude Lajoie ^{1,2}

¹ Department of Physical Activity Sciences, Université du Québec à Trois-Rivières, Trois-Rivières, QC, Canada

² Laboratoire de technologies & d'innovation pour la performance sportive (L-Tips), Université du Québec à Trois-Rivières, Trois-Rivières, QC, Canada

³ Department of Electrical and Computer Engineering, Université du Québec à Trois-Rivières, Trois-Rivières, QC, Canada

* Correspondence: (JCF) jules.cusson.fradet@uqtr.ca

Received: 11 March 2023; Accepted: 20 March 2023; Published: 23 October 2023

Keywords: Validity; Reliability; Endurance Training

1. Introduction

Monitoring the cyclists' responses to training is important for optimizing training and improving cycling performance (Galán-Rioja, Gonzalez-Ravé, González-Mohino, & Seiler, 2023; Jeffries et al., 2021). However, this process can be costly and time-consuming. Submaximal fitness tests provide a practical alternative to traditional maximal tests, allowing the monitoring of training effects while minimizing disruptions to the athlete's training and competition schedule (Shushan et al., 2022).

Recently, Sangan, Hopker, Davison, & McLaren (2021) proposed the self-paced submaximal run test (SRT_{RPE}) as a more practical form of the running version of the Lamberts and Lambert submaximal cycling test (LSCT) (Lamberts, Swart, Noakes, & Lambert, 2011). The LCST measures cycling power output (w) or running velocity (v) and ratings of perceived effort (RPE) for two 6-minute stages and one 3-minute stage of increasing intensity, which are determined by a fixed percentage of the maximum heart rate (Lamberts et al., 2011; Vesterinen et al., 2016). In contrast, the SRT_{RPE} monitors v and heart rate (HR_{ex}) during three 3-minute stages prescribed by RPE 10, 13, and 17 (Sangan et al., 2021).

The self-paced nature of the SRT_{RPE} makes it potentially more specific to the

spacing demands of competition and, more importantly, less cumbersome since there is no requirement to collect and to store RPE data for later analysis, nor to do a prior maximal testing to establish the relative intensity of each stage. Furthermore, standardizing intensity by RPE could provide a better insight into the individual's responses to endurance training by reflecting more accurately individual differences in the exercise intensity domains (Iannetta et al., 2020; Seip, Snead, Pierce, Stein, & Weltman, 1991). Sangan et al. (2021) reported promising findings showing that v , at all stages of the SRT_{RPE}, was largely correlated with aerobic fitness parameters such as maximal oxygen uptake ($\dot{V}O_{2max}$), v at $\dot{V}O_{2max}$ ($v\dot{V}O_{2max}$), and v at lactate threshold 2 ($vLT2$). Moreover, test-retest analysis showed acceptable reliability with coefficients of variation between 2.5% and 5.6% for v and HR_{ex} at all stages (Sangan et al., 2021).

Given the supportive findings and potential benefits of a self-paced design, the aim of our study was to evaluate the feasibility of a cycling-specific adaptation of the SRT_{RPE} protocol. The modified protocol incorporates the latest research on RPE and is tailored towards the specific requirements of road cycling (Ebert, Martin, Stephens, & Withers, 2006; Lopes, Pereira, & Silva, 2022). The self-paced submaximal cycling test



(SCT_{RPE}) measures w and HR_{ex} responses during three 2-minute stages using Borg's CR100 scale (RPE Moderate, Hard, Very Hard) and evaluates neuromuscular function with one second peak power (w_{sprint}) and post-exercise heart rate recovery (HRR).

2. Materials and Methods

Twenty-one trained to highly trained (27 ± 6 y, 71.8 ± 9.6 kg, $\dot{V}O_{2\text{peak}} 66 \pm 7$ ml·kg⁻¹·min⁻¹) endurance cyclists (including 3 females) received standardized RPE instructions and anchoring procedures. Then, they underwent the first trial of the SCT_{RPE}. They also performed three 6-second maximal sprints and a graded exercise test (GXT) to determine their maximal neuromuscular one second peak power ($w_{\text{MAXsprint}}$), ventilatory thresholds (VT1, VT2), peak oxygen consumption ($\dot{V}O_{2\text{peak}}$), and peak power (w_{peak}). Within a period of 2 to 14 days, the participants completed two additional trials of the SCT_{RPE} to evaluate the test-retest reliability of HRR, w_{sprint} , HR_{ex}, and w at each stage of the test. Correlations were used to examine the relationship between the SCT_{RPE} and the parameters measured during the GXT to establish convergent validity.

3. Results

Power output measured during all stages of the SCT_{RPE} had moderate to very large correlations with the parameters measured during the GXT, ranging from .33-.84. Notably, the highest correlation ($r = .84$) was observed between w_{peak} and w at RPE Very Hard (Figure 2). The intraclass correlation coefficients (ICC_{3,1}) for w and HR_{ex} at all stages of the SCT_{RPE} ranged from .79-.87, indicating high reliability, with typical errors (TEs) ranging from 2.7% to 7.7% (Table 1). The reliability of HRR expressed as a percentage of the peak heart rate recorded during the trial (HRR_{%hrpeak}) and w_{sprint} was found to be high and very high, respectively, with ICC_{3,1} value of .76 and .91, and TEs of 4% and 7% (Table 1).




Figure 1. Box plots showing the differences in power output at each stage of the SCT_{RPE} over 3 repeated trials. The box defines the upper and lower quartiles and the median. Whiskers show the minimum and maximum differences. RPE indicates rating of perceived effort; SCT_{RPE}, self-paced submaximal cycling test.

Table 1. Test-Retest Reliability of the Parameters of the Self-Paced Submaximal Cycling Test, Over 3 Repeated Trials (n = 21).

Parameter	Mean (SD)	TE	CV (%)	ICC
Power output (W)				
RPE Moderate	205.6 (34.47)	15.83	7.7	0.79
RPE Hard	284.81 (41.54)	17.37	6.1	0.83
RPE Very Hard	340.59 (54.44)	19.97	5.86	0.87
Heart rate (beats/min)				
RPE Moderate	143.17 (14.62)	6.31	4.41	0.8
RPE Hard	161.36 (13.84)	5.91	3.66	0.81
RPE Very Hard	175.58 (12.8)	4.78	2.72	0.86
Others				
HRR (beats/min)	44.51 (13.76)	6.17	13.86	0.75
HRR _{%hrpeak} (%)	25.38 (8.12)	4.05	-	0.76
w_{sprint} (W)	1147.03 (263.94)	81.2	7.08	0.91

Abbreviations: HRR, heart rate recovery; HRR_{%hrpeak}, heart rate recovery expressed as a percentage of the peak heart rate recorded during the trial; w_{sprint} , neuromuscular function; TE, Typical error; CV (%), TE expressed as a coefficient of variation; ICC, Intraclass correlation coefficients₃₋₁.




Figure 2. Pearson correlation (r) with 95% confidence interval of power output at RPE Very Hard with w_{peak} from the graded exercise test.

4. Conclusions

The study findings suggest that the SCT_{RPE} is a reliable and valid measure of endurance cycling fitness. The high intraclass correlation coefficients and good convergent validity with the parameters of the GXT, particularly W_{peak} , indicate that the SCT_{RPE} could be an effective and practical test to monitor endurance cycling athletes' responses to training.

5. Practical Applications

Athletes and coaches using the SCT_{RPE} may be able to observe changes in training status when the power output exceeds the TE measured for each stage. For example, if the power output of a cyclist during the last stage (RPE Very Hard) of the SCT_{RPE} exceeds by 21W his power output of the previous test, we can assume that his cycling fitness is rising because the change is greater than the TE of 20W for this stage.

Funding: This research received no external funding.

Acknowledgments: The authors would like to thank all participants for their voluntary participation.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Ebert, T. R., Martin, D. T., Stephens, B., & Withers, R. T. (2006). Power output during a professional men's road-cycling tour. *International Journal of Sports Physiology and Performance*, 1(4), 324–335. <https://doi.org/10.1123/ijspp.1.4.324>
- Galán-Rioja, M. Á., Gonzalez-Ravé, J. M., González-Mohino, F., & Seiler, S. (2023). Training Periodization, Intensity Distribution, and Volume in Trained Cyclists: A Systematic Review. *International Journal of Sports Physiology and Performance*, 1–11. <https://doi.org/10.1123/ijspp.2022-0302>
- Iannetta, D., Inglis, E. C., Mattu, A. T., Fontana, F. Y., Pogliaghi, S., Keir, D. A., & Murias, J. M. (2020). A Critical Evaluation of Current Methods for Exercise Prescription in Women and Men. *Medicine and Science in Sports and Exercise*, 52(2), 466–473. <https://doi.org/10.1249/MSS.0000000000002147>
- Jeffries, A. C., Marcora, S. M., Coutts, A. J., Wallace, L., McCall, A., & Impellizzeri, F. M. (2021). Development of a Revised Conceptual Framework of Physical Training for Use in Research and Practice. *Sports Medicine*. <https://doi.org/10.1007/s40279-021-01551-5>
- Lamberts, R. P., Swart, J., Noakes, T. D., & Lambert, M. I. (2011). A novel submaximal cycle test to monitor fatigue and predict cycling performance. *British Journal of Sports Medicine*, 45(10), 797–804. <https://doi.org/10.1136/bjsm.2009.061325>
- Lopes, T. R., Pereira, H. M., & Silva, B. M. (2022). Perceived Exertion: Revisiting the History and Updating the Neurophysiology and the Practical Applications. *International Journal of Environmental Research and Public Health*, 19(21). <https://doi.org/10.3390/ijerph192114439>
- Sangan, H. F., Hopker, J. G., Davison, G., & McLaren, S. J. (2021). The Self-Paced Submaximal Run Test: Associations With the Graded Exercise Test and Reliability. *International Journal of Sports Physiology and Performance*, 1(aop), 1–9. <https://doi.org/10.1123/ijspp.2020-0904>
- Seip, R. L., Snead, D., Pierce, E. F., Stein, P., & Weltman, A. (1991). Perceptual responses and blood lactate concentration: effect of training state. *Medicine and Science in Sports and Exercise*, 23(1), 80–87. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/1997816>
- Shushan, T., McLaren, S. J., Buchheit, M., Scott, T. J., Barrett, S., & Lovell, R. (2022). Submaximal Fitness Tests in Team Sports: A Theoretical Framework for Evaluating Physiological State. *Sports Medicine*, 52(11), 2605–2626. <https://doi.org/10.1007/s40279-022-01712-0>
- Vesterinen, V., Nummela, A., Äyrämö, S., Laine, T., Hynynen, E., Mikkola, J., & Häkkinen, K. (2016). Monitoring Training Adaptation With a Submaximal Running Test Under Field Conditions. *International Journal of Sports Physiology and Performance*, 11(3), 393–399. <https://doi.org/10.1123/ijspp.2015-0366>